



Review

Capnocytophaga canimorsus

Wim Gaastra ^a, Len J.A. Lipman ^{b,*}

^a Department of Infectious Diseases and Immunology, Faculty of Veterinary Medicine, Utrecht University, The Netherlands

^b Division of Public Health and Food Safety, Institute for Risk Assessment Sciences, Utrecht University, Yalelaan 2, 3508 TD Utrecht, The Netherlands

ARTICLE INFO

Article history:

Received 5 November 2008

Received in revised form 22 January 2009

Accepted 27 January 2009

Keywords:

Capnocytophaga canimorsus

Zoonosis

Sepsis

Virulence

Risk factors

Therapy

ABSTRACT

Capnocytophaga canimorsus is a commensal bacterium in the oral flora of dogs and cats. The bacterium is a zoonotic agent and has been isolated from humans, infected by dog or cat bites, scratches, licks or simply exposure to dogs or cats. Here the infectious agent, its pathogenicity and potential virulence factors, infection in animals and humans, diagnostic methods, prevalence, therapy and prevention are described. Suggestions for future research are given.

© 2009 Elsevier B.V. All rights reserved.

Contents

1. Introduction	339
2. The infectious agent	340
2.1. Historical perspective.	340
2.2. Phenotypic and genotypic characteristics	340
2.3. Cultural properties	340
2.4. Biochemical characteristics	341
2.5. Pathogenicity and virulence factors.	341
3. Infection in animals.	341
4. Human infections and clinical symptoms.	342
5. Risk groups.	342
6. Diagnostics.	343
7. Therapy and prevention	343
8. Prevalence	344
9. Conclusions	344
10. Future research	345
Acknowledgements.	345
References	345

1. Introduction

Worldwide millions of people are bitten by animals each year. It is estimated that 50% of all Americans will be

* Corresponding author. Tel.: +31 302535342.
E-mail address: L.J.A.Lipman@iras.uu.nl (Len J.A. Lipman).

bitten at least once in their lifetime. Ninety percent of these bites are caused by dogs and cats. The majority of dog and cat bite wounds is minor and no medical attention is sought by the victim. Occasionally a dog or cat bite is complicated and an overwhelming systemic infection occurs. In the USA, the annual mortality rate due to dog and cat bites is 6.7 per 10⁸ persons, albeit that not all these fatalities are caused by infections (Griego et al., 1995).

The genus *Capnocytophaga* consists of seven species which inhabit the oral cavity of humans or domestic animals. *Capnocytophaga gingivalis*, *Capnocytophaga ochracea*, *Capnocytophaga sputigena*, *Capnocytophaga granulosa* and *Capnocytophaga haemolytica* are inhabitants of the human oral cavity and have been associated with periodontitis. *Capnocytophaga cynodegmi* and *Capnocytophaga canimorsus* (latin canis = dog and latin morsus = bite) are part of the oral microbiota of canines and more rarely of cats. Eight percent of 50 dogs tested positive for *C. canimorsus* in the first report on the prevalence of *C. canimorsus* (Bailie et al., 1978). In a later report *C. canimorsus* could be cultured from 26% of the dogs tested and from 18% of the tested cats. The agent could not be cultured from hamsters and humans (Blanche et al., 1998). In our laboratory, a PCR test with primers for the 16S rRNA gene of *C. canimorsus* was positive for 41% of the dogs (de Poel et al., unpublished results). An even higher percentage was observed with a similar PCR by van Dam et al. (unpublished results). In a single report, the isolation of *C. canimorsus* (DF-2) from sheep and cattle (25–30% of the animals tested) but not from pigs was reported (Westwell et al., 1989). To our knowledge no attempt to confirm this observation has ever been made. *C. canimorsus* can cause zoonotic infections ranging from very mild flu like symptoms to fatal sepsis (de Boer et al., 2007; Pers et al., 1996). *C. canimorsus* infections are associated with dog and cat bites (54% of cases), dog and cat scratches (8.5% of cases) or close animal contact (27% of cases), such as licking of human wounds (Lion et al., 1996). Consequently human-to-human transmission of *C. canimorsus* has not been reported apart from one case where it could not be excluded 100% (Risi and Spangler, 2006). Studies on *C. canimorsus* and *C. cynodegmi* infections have mainly been initiated from an interest in human medicine for rapid diagnosis and antibiotic treatment, prompted by the often dramatic outcome of an infection with *C. canimorsus*. Since its first description in 1976 approximately 200 human cases of *C. canimorsus* infection have been reported worldwide (Macrea et al., 2008). Surprisingly little research on this zoonotic agent has been performed from a veterinary background. Regarding the millennia old bond between men and dog, this infection is probably not new, but only now it can be recognized due to the advances in modern medicine over the last four decades.

2. The infectious agent

2.1. Historical perspective

In 1976, a patient with meningitis and sepsis, caused by an unidentified Gram-negative bacillus, after a recent dog bite was first described (Bobo and Newton, 1976). At that

time, the phenotypic as well as the biochemical characteristics of the agent also became known. We now know that this Gram-negative bacillus probably was a member of the *Capnocytophaga* family, very likely *C. canimorsus* (CDC group DF-2 or dysgonic fermentor 2, dysgonic = slow and relatively poor growth of a bacterial culture). One year later, 17 cases of infection after recent dog bites, the majority caused by an unidentified Gram-negative rod were described (Butler et al., 1977). The nature of the infection varied from cellulitis to meningitis, endocarditis to organ failure and fatal sepsis. The organism was given its current name in 1989, based on its CO₂ requirement and usual mode of transmission (Brenner et al., 1989). Since then, *C. canimorsus* infections were found to occur worldwide and have been reported from the United States, Canada, Europe, Australia and South Africa.

2.2. Phenotypic and genotypic characteristics

C. canimorsus are capnophilic (greek:carbon dioxide loving) facultative anaerobic, fastidious Gram-negative rods (1–4 µm long, Fig. 1). They are fusiform or filamentous gliding bacteria and closely related to *Fusobacterium* and *Bacteroides* species (Andersen and Pedersen, 1992; Deshmukh et al., 2004). However, the gliding motility is not always apparent. The G+C content of *C. canimorsus* DNA is 35% (Speck et al., 1987; Brenner et al., 1989).

2.3. Cultural properties

Classically, the organism was regarded difficult to culture because of its specific requirements for nutrients and thus may have easily escaped detection. *C. canimorsus* has been shown to require large amounts of exogenous iron for growth (Brenner et al., 1989). It grows slowly on blood (5% sheep blood in Columbia agar) or chocolate agar in 10% CO₂ (Ciantar et al., 2001). Media had to be incubated at 37 °C for at least 5 days. Colonies could be not visible for 2–7 days, on blood agar. Initially they are pinpoint in size and later become larger, convex and smooth. Colonies are non hemolytic with the characteristic shiny spreading edge and finger-like projections of bacteria with gliding motility (Andersen and Pedersen, 1992; Deshmukh et al.,




Fig. 1. Gram stain of *C. canimorsus* isolate from a rabbit bitten by a dog.

2004). Although the color of colonies may vary from pink to yellow, the colonial mass of all isolates is yellow when scraped from the agar plate (Forlenza, 1991). Horse blood agar plates were found to be superior to sheep blood but do not seem to be used often (Westwell et al., 1989; Pers et al., 1996). The organism does not grow on MacConkey agar. Recently, the growth medium was optimized (Heart Infusion agar Difco with 5% sheep blood) and the bacteria were shown to grow in 2 days at 37 °C in 5% CO₂ (Shin et al., 2007; Mally et al., 2008). To our knowledge, these growth conditions have only been used under experimental conditions and not yet in a clinical setting. It is to be expected that faster growth of *C. canimorsus* will be reflected in the speed with which diagnostics can be performed and therefore lead to less fatalities due to this organism.

2.4. Biochemical characteristics

Biochemical tests are difficult to perform due to the slow growth of the organism. It is catalase, oxidase, ONPG (*O*-nitrophenyl-β-D-galactoside) and arginine dihydrolase positive and negative for urease, nitrate, indole, DNase, gelatin, lysine and ornithine (Hicklin et al., 1987; Brenner et al., 1989). Members of the genus *Capnocytophaga* can use various carbohydrates like glucose, dextran, glycogen, inulin or starch as fermentable substrates and energy source. For a comparison of the biochemical characteristics of *C. ochracea*, *sputigena*, *gingivalis*, *canimorsus* and *cynodegmi*, see Forlenza (1991). Gas–liquid chromatography of cellular fatty acids provides an additional rapid test for differentiation of Gram-negative bacteria associated with dog-bite infections, including *C. canimorsus* (Dees et al., 1981).

2.5. Pathogenicity and virulence factors

Little is known about the pathogenesis of *C. canimorsus* infection. The bacterium was reported to multiply in mouse macrophages and to be cytotoxic probably because it produces a toxin (Fischer et al., 1995). In vitro, a very low level of cytokine production i.e. in comparison to *N. meningitidis* was observed (Frieling et al., 1997). Cytotoxicity could not be demonstrated by Shin et al. (2007) in 10 strains of *C. canimorsus* tested, including the same strain as in the original study. These strains also did not affect the viability of macrophages from different origin (mouse and human). The clinical course of *C. canimorsus* infection strongly indicates that the organism is able to avoid the immune system (at least in the early stages of infection). The absence of a proinflammatory response is due to the fact that *C. canimorsus* does not interact with human Toll-like receptor 4 (TLR4). The effector for TLR4 is the lipid A moiety of LPS, which indicates that lipid A of *C. canimorsus* is different from that of other pathogens. *C. canimorsus* is able to down regulate TLR4 and the proinflammatory signaling cascade (Shin et al., 2007). Not only is *C. canimorsus* resistant against phagocytosis and killing, it also blocks the killing of unrelated bacteria by macrophages (Meyer et al., 2008). Despite being classified as a fastidious grower, *C. canimorsus* exhibits robust growth

when it is in direct contact with mammalian cells, including phagocytes (Mally et al., 2008). This is dependent on a surface located sialidase, which allows the bacterium to use internal amino-sugars from glycan chains of host cell glycoproteins (Mally et al., 2008). The molecular biology of *C. canimorsus* is still in its infancy, but promising genetic tools for this bacterium have recently been developed using a naturally occurring plasmid found in one of the eight strains tested (Mally and Cornelis, 2008).

3. Infection in animals

The clinical syndrome in rabbits after experimental infection with *C. canimorsus* was characterized by disseminated intravascular coagulation, cellular necrosis in organs like kidneys and adrenal glands, cutaneous gangrene, thrombocytopenia, hypotension, hemorrhagic diathesis with purpuric lesions and petachiae and renal failure and is similar to that observed in man (Piccininno et al., 1984).

Only two reports have appeared in species other than humans that suffered from a dog inflicted *C. canimorsus* infection. In one case the bacterium was cultured from an infected, bite wound in a pet rabbit (van Duijkeren et al., 2006) (Fig. 2). In the other *C. canimorsus* was isolated from an infected bite wound in a dog (Meyers et al., 2008). The dog did not have bacteremia, but *C. canimorsus* could still be isolated from the wound 2 days after the primary isolation (J. Picard, personal communication). Curiously both bite wounds were on the head. No *C. canimorsus* infections of cats after a bite incidence have been reported, but there is a case report of a cat with chronic sinusitis and rhinitis due to a *Capnocytophaga* infection. The organism was identified by a 97% identity with *C. canimorsus* of its 16S rDNA sequence (Frey et al., 2003). The incidence of bite wounds inflicted in dogs and cats by dogs or cats is not well known. It is estimated that they comprise between 10 and 15% of all trauma related emergencies (Kolata et al., 1974). Like with human bite wounds many wounds are not reported and owners do not seek veterinary attention. In the mobile, elastic skin of dogs and cats, bite injuries may be less noticeable or may migrate through deeper tissues and escape detection. Dogs and cats may also be less prone



Fig. 2. Rabbit with *C. canimorsus* infection after a dog bite in the head.

to *C. canimorsus* infection, which might explain the very low number of reported *C. canimorsus* infection in animals.

4. Human infections and clinical symptoms

The incubation period from the dog bite to the onset of systemic symptoms is about 5 days. On average 7 days pass between the time of the bite and hospitalization (range 1–14 days) (LeMoal et al., 2003). Patients who arrive at the emergency room within 8–12 h after a dog bite may show local lesions without significant signs of inflammation. At later stages, the infection may have symptoms as: localized cellulitis, pain at the site of injury, a purulent discharge, lymphangitis and regional lymphadenopathy. The initial symptoms of septicemia are fever (78% of patients), chills (46%), myalgia (31%), vomiting (31%), diarrhea (26%), abdominal pain (26%), malaise (26%), dyspnea (23%), mental confusion (23%) and headache (18%) (Pers et al., 1996). A fulminant and severe course of the infection may occur in immunocompromised persons, characterized by sepsis, meningitis, osteomyelitis, peritonitis, endocarditis, pneumonia, purulent arthritis, disseminated intravascular coagulation (DIC) and fulminant purpura but has also been observed in previously healthy persons (Hantson et al., 1991). DIC can lead to peripheral gangrene of the upper and lower extremities, lips nose and ears (often resulting in amputation). Purpura is a group of blood disorders that cause bleeding in the tissue from the small blood vessels, shock and multi-organ failure. They are often under or in the skin as tiny pinpoint petechiae (point-shaped skin bleedings or larger blue spots (ecchymosis). Hemolytic uremic syndrome and Waterhouse–Friderichsen syndrome have also been reported (Tobé et al., 1999; Mirza et al., 2000; Mulder et al., 2001; Macrea et al., 2008). *C. canimorsus* septicemia has been associated with up to 30% mortality (Lion et al., 1996) and significant morbidity, including amputation secondary to gangrene, myocardial infarction and renal failure. Curiously, Lion et al. (1996) report 22% mortality in asplenic individuals, 28% in individuals with other predisposing conditions (alcohol abuse, immunosuppression) and 32% in individuals with no risk factor. The mortality rate from *C. canimorsus* meningitis is low (5%) compared to that from *C. canimorsus* septicemia (LeMoal et al., 2003; Janda et al., 2006). In a review of 12 cases of *C. canimorsus* endocarditis a mortality rate of 25% was noted (Sandoe, 2004). A fatal outcome of the infection in non-immunocompromised individuals occurs with lower frequency than in immunocompromised persons. Four of these cases were reported until 1991 and several others between 1991 and 2008 (Hicklin et al., 1987; Hantson et al., 1991; Deshmukh et al., 2004). Ocular infections associated with cat scratches or cat bites have been reported several times (Glasser, 1986; Zimmer-Gallen and Pach, 1996; Papadaki et al., 2008).

5. Risk groups

At special risk are animal keepers, veterinarians, breeders and pet owners. A survey of veterinarians in the US showed that 65% sustained a major animal related injury of which animal bites and scratches accounted for

38% of the traumas. Dogs and cats were involved in 24% and 10% of the injuries. In their entire careers, 92% of the veterinarians were bitten by a dog, 81% by a cat and 72% were scratched by a cat (Landercasper et al., 1988). Yet only few reports of persons in this risk group infected with *C. canimorsus* are available: an employee in an animal shelter (Deshmukh et al., 2004), a pet shop worker (Hantson et al., 1991), three veterinarians (Job et al., 1988; de Smet et al., 1990; Chodosh, 2001), a kennel operator (Tison and Latimer, 1986.) and a veterinary worker (Bobo and Newton, 1976), which is much lower than for rat bite fever, for which the same group is also at risk (Gastra et al., 2009). Two cases in veterinarians were not caused by bites or scratches, but by fracture of a tooth during manual extraction which struck the veterinarians in the eye (de Smet et al., 1990; Chodosh, 2001). In one case it was the tooth of a cat, in the other of a dog. *C. canimorsus* is primarily an opportunistic pathogen that infects individuals weakened by a compromised immunity. Most infections in humans occur among people older than 40 years, but infections have been reported in a 4-month-old and 83 year-old patient, the ratio male to female is 3:1 (Danker et al., 1987; Pers et al., 1996; Deshmukh et al., 2004; Sandoe, 2004; Janda et al., 2006). Sixty percent of the people that get an infection with this organism have predisposing conditions. Those who have had a splenectomy (33% of cases) comprise a high-risk group with a 30–60 times increased risk of fatal septicemia (Singer, 1973, Deprés-Brummer et al., 2001). Often, already within 24 h of symptom onset, overwhelming postsplenectomy infections (OPSI) can result in a rapidly deteriorating clinical course that progresses to multiple organ system failure and death (Band et al., 2008; Sawmiller et al., 1998). In the United States each year 25,000 splenectomies are performed (Sumaraju et al., 2001). Together with the thousands of functionally hyposplenic or asplenic patients a significant number of people are therefore at marked risk for OPSI, which has an estimated mortality rate of 70% (Brigden and Pattullo, 1999). Asplenic individuals suffer deficient IgM and IgG production and delayed macrophage mobilization. They also produce less tuftsin, a protein derived from IgG that stimulates phagocytosis (August, 1988). Liver disease caused by alcohol abuse (24% of cases) and the use of cortisone (3% of cases) are other predisposing factors for the infection (Lion et al., 1996). Hematologic malignancy, neutropenia and chronic lung disease were also considered as predisposing conditions (Dire et al., 1994). Up to 40% of cases however occur in previously healthy adults (LeMoal et al., 2003). Alcohol abuse, asplenia, hemochromatosis, β -thalassemia major and tobacco smoking lead to absorption of twice the amount of alimentary iron as in normal persons. In two cases of immune-competent persons with *C. canimorsus* infection, both women, heavy smoking was explicitly mentioned (Finn et al., 1996; Aslam, 1999). An alternative explanation for why these conditions form a risk for *C. canimorsus* infection might be that this two-fold increase in serum iron values much better supports the growth of the poor iron scavenger *C. canimorsus* (Weinberg, 2000). It was suggested that the presence of detectable amounts of *C. canimorsus* in only one-quarter of dogs and cats might be

due to the fact that positive animals have insufficient amounts of lactoferrin and other iron binding factors in their oral cavity (Weinberg, 2000). Since the 25% is likely to be an underestimation based on data with improved selective cultivation conditions (see above) and PCR (see above), the suggested lack of lactoferrin in *C. canimorsus* positive animals is unlikely. However, iron overload was suggested as a risk factor in a case of *C. canimorsus* infection in a patient receiving numerous blood transfusions over a 2-year period (Pletschette et al., 1992).

6. Diagnostics

The diagnosis *C. canimorsus* infection is usually made based on the bacterial culture of blood (88% of cases), other body fluids (cerebrospinal fluid, 7% of cases) or less frequently from the bite wound or tissue from the bitten individual (Janda et al., 2006). Polyanethole-sulfonate, an anticoagulant frequently present in automatic blood culture systems, inhibits the growth of *C. canimorsus* (Sowden et al., 1995). Swabs of the mouth of the dog that bit the patients have been taken in a number of cases, but growth of *C. canimorsus* was only obtained in half of them (Martone et al., 1980; Howell and Woodward, 1990; Hantson et al., 1991; Valtonen et al., 1995; Finn et al., 1996; Phipps et al., 2002; van de Ven et al., 2004). An indication of a *C. canimorsus* infection can already be obtained when a Gram stain of the buffy coat of peripheral blood shows many extracellular Gram-negative rods and several intracellular copies of the same rods in almost all neutrophils (Pedersen et al., 1993; Sawmiller et al., 1998; Mirza et al., 2000). In cases of overwhelming *C. canimorsus* septicemia, whole blood smears have also been reported positive for Gram-negative rods in the Gram stain (Holmes and Kozinn, 1986; Ndon, 1992; Newton, 2006; Kleijnen-Grebien et al., 2008). Examination of the buffy coat by Gram stain in splenectomized patients is particularly useful when *C. canimorsus* infection is suspected, since several days may pass before the organism is identified in the laboratory (Hicklin et al., 1987). The slow growth of *C. canimorsus* may lead to diagnostic problems since culture plates are routinely discarded after 5 days and some strains do not grow at all. A Gram stain of the cerebrospinal fluid showed Gram-negative bacilli in 65% of 20 cases of *C. canimorsus* meningitis (de Boer et al., 2007).

The diagnosis of *C. canimorsus* infection was evaluated by Janda et al. (2006) with isolates collected over a 30-year period. It was concluded that many laboratories are unable to identify *C. canimorsus* isolates and report them as (fastidious) Gram-negative rods. In their identification procedure the authors use a combination of biochemical tests, fatty acid methyl ester analysis and 16S rDNA sequencing (Janda et al., 2006). The number of cases in which broad range PCR followed by DNA sequencing is used for diagnostic purposes is increasing (Ciantar et al., 2005; Gottwein et al., 2006; Janda et al., 2006; Meybeck et al., 2006; Wareham et al., 2007; Le Meur et al., 2008; Kleijnen-Grebien et al., 2008; Papadaki et al., 2008). It should be noted that in one case, amplification of *C. canimorsus* DNA from an infected valve was obtained while

attempts to amplify the DNA from the positive blood culture of the same patient was unsuccessful, probably due to the presence of polyanethylsulfonate which inhibits the PCR reaction (Wareham et al., 2007).

7. Therapy and prevention

The first choice antibiotic for infection with *C. canimorsus* is penicillin G, although resistance of isolates has been mentioned (Meybeck et al., 2006). After investigation of the available literature we came to the same conclusion as LeMoal et al. (2003), that no β -lactamase producing *C. canimorsus* strain has been reported yet. There are however, several reports on β -lactamase producing *Capnocytophaga* species (Arlet et al., 1987; Roscoe et al., 1992; Jolivet-Gougeon et al., 2007). With broth or agar dilution techniques, susceptibility for penicillins, imipenem, clindamycin, chloramphenicol, third generation cephalosporins, fluoroquinolones, erythromycin, doxycycline and metranidazole has been demonstrated. Aztronam, polymyxin, fosfomycin, trimethoprim/sulfamethoxazole and aminoglycosides are not effective against this bacterium (Verghese et al., 1988; Bremmelgaard et al., 1989; Deshmukh et al., 2004). However, in the literature disagreement on the results for trimethoprim/sulfamethoxazole, aminoglycosides and vancomycin can be found, which seem to depend on the method used. Routine prophylaxis with amoxicillin-clavulanic acid has been proposed for all immunocompromised patients and in cases seen within 8 h after moderate to severe crush injuries or when the bones and joints are involved (Goldstein, 2005). Immediate cleaning of the wound and disposal of scrap fabric filler are important during the antibiotic therapy. Although systemic administration of antimicrobial agents is controversial for healthy persons, those from whom the spleen was removed should be given prophylactic antibiotics against this organism after a dog bite or in the case of an open wound which has been in contact with the saliva of a dog. In several countries systemic antibiotic treatment is recommended for all persons (not only immunocompromised patients) after a dog bite. The clinical course of a patient with *C. canimorsus* sepsis, which presented as an acute abdomen described by Sawmiller et al. (1998) clearly demonstrates the importance of early diagnosis and treatment of an overwhelming postsplenectomy infection. One may wonder whether there is an indication to test healthy pets for *C. canimorsus*, in households with high-risk individuals and if there is an indication to treat colonized pets in these households. Given the high prevalence of dogs when tested by PCR for *C. canimorsus*, it is very likely that dogs in high-risk households will test positive. It is unclear what the advance of treating the dogs in these households will be since a treated “*C. canimorsus* free” dog can readily be colonized again after contact with other dogs. There are no reports on dogs that have been tested several times over a period of time. So it is not known whether a negative dog will stay negative during its whole life or whether a positive dog will always test positive a next time. Furthermore, since a PCR test for *C. canimorsus* has not been validated it is not clear what the value of a

negative PCR is with respect to that particular dog being colonized or not

8. Prevalence

Worldwide millions of people are bitten by animals each year (one out of every two persons will be bitten by an animal at some point in his lifetime (Greene and Goldstein, 2006). Ninety percent of the bites are by dogs and cats (Griego et al., 1995; Matter and Sentinella, 1998). Less than half of the bites are reported and 18% of these persons seek medical attention (Overall and Love, 2001). Most bites are minor and occur (in descending order of incidence) on the extremities (hands and feet), head and neck and the fuselage. Although 80% of dog bite wounds inflicted on people culture positive for bacteria, the chance that a dog bite gets infected is between 3 and 20% (Underman, 1987; Goldstein, 1992). For cat bites, this is between 20 and 50% (Talan et al., 1999). Usually the infection is caused by a combination of aerobic and anaerobic bacteria. In the Netherlands yearly between 50,000 and 100,000 people get bitten by a pet animal. The number of registered bite incidents by dogs or cats in children under the age of 10 in The Netherlands is about 4800 per year (Deprés-Brummer et al., 2001). Since 1976, worldwide approximately 200 cases of septicemia, meningitis or gangrene caused by *C. canimorsus* have been reported in humans bitten, scratched or licked by a dog or cat (Brenner et al., 1989; Blanche et al., 1998; Tierney et al., 2006). Even though almost equal percentages of dogs and cats (24% vs. 18%) (Blanche et al., 1998) carry *C. canimorsus* in the oral cavity, the majority of cases is caused by dog bites or dog contact. Cat bites, scratches or contact with cats has been reported in less than 10% of the cases. The deep, sharp, needle like punctures with relatively little tissue damage through cat bites are probably creating less favorable conditions for the growth of the bacterium than the larger superficial wounds with a lot of tissue damage caused by dog bites (Carpenter et al., 1987; Scarlett et al., 1991; Mahrer and Raik, 1992; Valtonen et al., 1995; Zimmer-Gallen and Pach, 1996; McLean et al., 2004; Le Meur et al., 2008). The risk of infection depends on adequate wound treatment and host factors like age, health status.

Infection with *C. canimorsus* is not a notifiable disease. The number of reports on *C. canimorsus* infection is increasing, but it is still a relatively rarely reported disease. The frequency of *C. canimorsus* infection is very likely underestimated (Janda et al., 2006). Presently, not even all severe and lethal infections are reported but only those cases where the clinical picture has not been seen before. Additionally, patients who are supplied with antibiotics in time do not develop a *C. canimorsus* infection. This may also influence the infection rate. The bacteria may be transmitted in various ways by close contact between pets and their owners. Transmission is not necessarily followed by multiplication of the bacteria in the human body (infection). Infection does not necessarily lead to serious clinical symptoms and humans with subclinical infections will not be reported to the physician. If the incubation period extends to several weeks and clinical symptoms are

aspecific, *C. canimorsus* infection is probably not considered when bites by dogs or cats or contact with these animals are not explicitly mentioned in the anamnesis.

Patients suspected of *C. canimorsus* infection will often be treated with antibiotics most of which will be active against the causative bacteria. The number of cases in which laboratory diagnostic examinations are carried out is therefore limited to very severe cases or when antibiotic therapy fails.

C. canimorsus may be difficult to grow in primary culture after antibiotic therapy and detection by broad range PCR and nucleotide sequence determination is operational in a limited number of diagnostic laboratories only. *C. canimorsus* strains may be misidentified despite the fact that the bacteriological characteristics are rather typical.

9. Conclusions

C. canimorsus is a commensal in the oral cavity of dogs and cats and is considered to be of low virulence under normal circumstances. The bacterium can cause severe illness in predisposed patients (i.e. those who have had a splenectomy, who are alcohol abusers, suffer from chronic lung or liver diseases or are on immunosuppressive medication). However, 40% of the reported cases occurred in previously healthy people.

The true number of *C. canimorsus* infections each year probably is largely underestimated due to the fastidious growth of the organism. The isolation from animal saliva is difficult and the organism is seldom isolated from bite wounds. The relatively more successful growth from blood culture can explain why the bacterium is only detected in cases of septicemia. Penicillin is considered the drug of choice in the case of *C. canimorsus* infection. The length of the treatment varies between reports (14–21 days). (Le Moal et al., 2002).

In the past, diagnosis and identification have been difficult due to the fastidious growth of the organism, which has led to fatal outcomes in cases where the patient probably could have survived since the bacterium is susceptible to a whole range of commonly used antibiotics. With the introduction of optimized growth conditions on selective media and PCR as a diagnostic tool this type of fatalities are expected to decrease. Because of the seriousness of *C. canimorsus* infection, taking this bacterium into account in the management of animal bites, is necessary. Moreover, high-risk patients should be informed about the risks of keeping pets. Not only do bites and scratches pose a risk to these patients, but other scenarios one not normally thinks of are possible. This can be illustrated by a case report where an 18-year-old person on automated peritoneal dialysis developed *C. canimorsus* peritonitis following the puncture of his dialysis tubing by a domestic cat that slept in his bed (Chadha and Warady, 1999).

Administration of antimicrobial agents is controversial for healthy persons, but it is justifiable for the following groups of bite victims: those with deep wounds, particularly to the face or hands and those at high risk for *C. canimorsus* infection e.g. immunocompromised patients.

10. Future research

More insight into the genetic properties and the molecular biology of *C. canimorsus* is a prerequisite for understanding the overwhelming sepsis, this organism can cause. In this respect whole genome sequencing of strains from different origin seems relevant. Questions to be answered are: are isolates from humans and dogs different? Or are the former a more virulent subset of the latter? Is there a difference between dog and cat isolates? Once the whole genome of *C. canimorsus* has been sequenced this might allow the identification of the genetic basis of virulence factors and the reason for the low virulence normally seen in immunocompetent bite victims. To improve diagnostics, media should be developed that allow rapid growth of the organism. For the following rapid identification the development of a species specific PCR is needed. This can eventually be followed by sequencing of the amplicon for confirmation. A systematic search for the presence of *C. canimorsus* in the oral cavity of other animals than cats and dogs is also needed to determine the potential risk with bites from these animals. So far there are no indications that point to a host specificity that is limited to dogs and cats.

Conflict of interest

None of the authors (Wim Gaastra, Len J.A. Lipman) has a financial or personal relationship with other people or organizations that could inappropriately influence or bias the paper entitled "*Capnocytophaga canimorsus*".

Acknowledgements

The authors are indebted to Chantal van Mourik, Engeline van Duijkeren (Veterinary Microbiological Diagnostics Centre, Veterinary Faculty, Utrecht University) and Marcel Leuven (Animal Clinic Leuven, Heerlen) for Figs. 1 and 2.

References

Andersen, H.K., Pedersen, M., 1992. Infective endocarditis with involvement of the tricuspid valve due to *Capnocytophaga canimorsus*. *Eur. J. Clin. Microbiol. Infect. Dis.* 11, 831–832.

Arlet, G., Sanson-Le Port, M.J., Casin, I.M., Ortenberg, M., Perol, Y., 1987. In vitro susceptibility of 96 *Capnocytophaga* strains, including a β -lactamase producer, to new β -lactam antibiotics and six quinolones. *Antimicrob. Agents Chemother.* 31, 1283–1284.

Aslam, A., 1999. Life-threatening *Capnocytophaga canimorsus* infection after dog bite. *J. R. Soc. Med.* 92, 140–141.

August, J.R., 1988. Dysgonic fermentor-2 infections. *J. Am. Vet. Med. Assoc.* 193, 1506–1508.

Bailie, W.E., Stowe, E.C., Schmitt, A.M., 1978. Aerobic bacterial flora of oral and nasal fluids of canines with reference to bacteria associated with bites. *J. Clin. Microbiol.* 7, 223–231.

Band, R.A., Gaieski, D.F., Goyal, M., Perrone, J., 2008. A 52-year-old man with malaise and petechial rash. *J. Emerg. Med.* (Epub ahead of publication).

Blanche, P., Bloch, E., Sicard, D., 1998. *Capnocytophaga canimorsus* in the oral flora of dogs and cats. *J. Infect.* 36, 134.

de Boer, M.G.J., Lambregts, P.C.L.A., van Dam, A.P., van't Wout, J.W., 2007. Meningitis caused by *Capnocytophaga canimorsus*: when to expect the unexpected. *Clin. Neurol. Neurosurg.* 109, 393–398.

Bobo, R.A., Newton, E.J., 1976. A previously undescribed Gram-negative bacillus causing septicemia and meningitis. *Am. J. Clin. Pathol.* 65, 564–569.

Bremmelgaard, A., Pers, C., Kristiansen, J.E., Korner, B., Heltberg, O., Frederiksen, W., 1989. Susceptibility testing of Danish isolates of *Capnocytophaga* and CDC group DF2 bacteria. *Acta Pathol. Microbiol. Immunol. Scand.* 97, 43–48.

Brenner, D.J., Hollis, D.G., Fanning, G.R., Weaver, R.E., 1989. *Capnocytophaga canimorsus* sp. Nov. (formerly CDC group DF-2), a cause of septicemia following dog bite, and *C. cynodegmi* sp. Nov., a cause of localized wound infection following dog bite. *J. Clin. Microbiol.* 27, 231–235.

Brigden, M.L., Pattullo, A.L., 1999. Prevention and management of overwhelming postsplenectomy infection—an update. *Crit. Care Med.* 27, 836–842.

Butler, T., Weaver, R.E., Ramani, T.K.V., Uyeda, C.T., Bobo, R.A., Ruy, J.S., Kohler, R.B., 1977. Unidentified gram-negative rod infection. *Ann. Intern. Med.* 86, 1–5.

Carpenter, P.D., Heppner, B.T., Gnann, J.W., 1987. DF-2 bacteremia following cat bites, report of two cases. *Am. J. Med.* 82, 621–623.

Chadha, V., Warady, B.A., 1999. *Capnocytophaga canimorsus* peritonitis in a pediatric peritoneal dialysis patient. *Pediatr. Nephrol.* 13, 646–648.

Chodosh, J., 2001. Cat's tooth keratitis: human corneal infection with *Capnocytophaga canimorsus*. *Cornea* 20, 661–663.

Ciantar, M., Spratt, D.A., Newman, H.N., Wilson, M., 2001. Assessment of five culture media for the growth and isolation of *Capnocytophaga* spp. *Clin. Microbiol. Infect.* 7, 158–160.

Ciantar, M., Newman, H.N., Wilson, M., Spratt, D.A., 2005. Molecular identification of *Capnocytophaga* spp. via 16S rRNA PCR-restriction fragment length polymorphism analysis. *J. Clin. Microbiol.* 43, 1894–1901.

Danker, W.M., Davis, C.E., Thompson, M.A., 1987. DF-2 bacteremia following dog bite in a 4 month old child. *Pediatr. Infect. Dis. J.* 6, 695–696.

Dees, S.B., Powell, J., Moss, C.W., Hollis, D.G., Weaver, R.E., 1981. Cellular fatty acid composition of organisms frequently associated with human infections resulting from dog bites: *Pasteurella multocida* and groups EF-4, Iij M-5 and DF-2. *J. Clin. Microbiol.* 14, 612–616.

Deprés-Brummer, P., Buijs, J., van Engelenburg, K.C.A., Oosten, H.R., 2001. *Capnocytophaga canimorsus* sepsis presenting as an acute abdomen in an aplenic patient. *Neth. J. Med.* 59, 213–217.

Deshmukh, P.M., Camp, C.J., Rose, F.B., Narayanan, S., 2004. *Capnocytophaga canimorsus* sepsis with purpura fulminans and symmetrical gangrene following a dog bite in a shelter employee. *Am. J. Med. Sci.* 327, 369–372.

Dire, D.J., Hogan, D.E., Riggs, M.W., 1994. A prospective evaluation of risk factors for infections from dog wounds. *Acad. Emerg. Med.* 1, 258–266.

van Duijkeren, E., van Mourik, C., Broekhuizen, M., Leuven, M., Gaastra, W., Houwers, D., 2006. First documented *Capnocytophaga canimorsus* infection in a species other than humans. *Vet. Microbiol.* 118, 148–150.

Finn, M., Dale, B., Isles, C., 1996. Beware of the dog! A syndrome resembling thrombotic thrombocytopenic purpura associated with *Capnocytophaga canimorsus* septicemia. *Nephrol. Dial. Transplant.* 11, 1839–1840.

Fischer, L.J., Weyant, R.S., White, E.H., Quinn, F.D., 1995. Intracellular multiplication and toxic destruction of cultured macrophages by *Capnocytophaga canimorsus*. *Infect. Immun.* 63, 3484–3490.

Forlenza, S.W., 1991. *Capnocytophaga*: an update. *Clin. Microbiol. Newslett.* 13, 89–91.

Frey, E., Pressler, B., Guy, J., Pitulle, C., Breitschwerdt, E.I., 2003. *Capnocytophaga* sp. isolated from a cat with chronic sinusitis and rhinitis. *J. Clin. Microbiol.* 41, 5321–5324.

Frieling, J.T.M., Mulder, J.A., Hendriks, T., Curfs, J.H.A.J., van der Linden, C.J., Sauerwein, R.W., 1997. Differential induction of pro- and anti-inflammatory cytokines in whole blood by bacteria: effects of antibiotic treatment. *Antimicrob. Agents Chemother.* 41, 1439–1443.

Gaastra, W., Boot, R., Ho, H.T.K., Lipman, L.J.A., 2009. Rat bite fever. *Vet. Microbiol.* 133, 211–228.

Glasser, D.B., 1986. Angular blepharitis by gram-negative bacillus DF-2. *Am. J. Ophthalmol.* 102, 119–120.

Goldstein, E.J.C., 1992. Bite wounds and infection. *Clin. Infect. Dis.* 14, 633–638.

Goldstein, E.F.C., 2005. Bites. In: Mandell, G.L., Bennett, J.E., Dolin, R. (Eds.), *Principles of Infectious Diseases*. 6th edition. Churchill Livingstone, New York, pp. 3552–3556.

Gottwein, J., Zbinden, R., Maibach, R.C., Herren, T., 2006. Etiologic diagnosis of *Capnocytophaga canimorsus* meningitis by broad range PCR. *Eur. J. Microbiol. Infect. Dis.* 25, 132–134.

Greene, C.E., Goldstein, E.J.C., 2006. Bite wound infections. In: Greene, C.E. (Ed.), *Infectious Diseases of the Dog and Cat*. 3th edition. Saunders, St. Louis, pp. 495–510.

Griego, R.D., Rosen, T., Orenge, I.F., Wolf, J.E., 1995. Dog, cat and human bites: a review. *J. Am. Acad. Dermatol.* 33, 1019–1029.

Hantson, P., Gautier, P.E., Vekemans, M.-E., Fievez, P., Evrard, P., Wauters, G., Mahieu, P., 1991. Fatal *Capnocytophaga canimorsus* septicemia in a previously healthy woman. *Ann. Emerg. Med.* 20, 93–94.

- Hicklin, H., Verghese, A., Alvarez, S., 1987. Dysgonic Fermentor 2 septicemia. *Rev. Infect. Dis.* 9, 884–890.
- Holmes, R.L., Kozinn, W.P., 1986. DF-2 septicemia following whirlpool spa immersion. *J. Clin. Microbiol.* 23, 627–628.
- Howell, J.M., Woodward, G.R., 1990. Precipitous hypotension in the emergency department caused by *Capnocytophaga canimorsus* sp. Nov. sepsis. *Am. J. Emerg. Med.* 8, 312–314.
- Janda, M.J., Graves, M.H., Lindquist, D., Probert, W.S., 2006. Diagnosing *Capnocytophaga canimorsus* infections. *Emerg. Infect. Dis.* 12, 340–342.
- Job, L., Horman, J.T., Grigor, J.K., Israel, E., 1988. Dysgonic fermenter-2: a clinico-epidemiologic review. *J. Emerg. Med.* 7, 185–192.
- Jolivet-Gougeon, A., Sixou, J.-L., Tamañal-Shacoori, Z., Bonnaure-Mallet, M., 2007. Antimicrobial treatment of *Capnocytophaga* infections. *Int. J. Antimicrob. Agents* 29, 367–373.
- Kolata, R.J., Kraut, N.H., Johnston, D.E., 1974. Patterns of trauma in urban dogs and cats: a study of 1000 cases. *J. Am. Vet. Med. Assoc.* 164, 499–502.
- Kleijnen-Greben, B., Boorsma, S., Stals, F.S., van Schelven, R., 2008. Fatale afloop van een sepsis met *Capnocytophaga canimorsus* na een triviale hondebeet. *Ned. Tijdschr. Geneesk.* 152, 1882–1885 (in Dutch).
- Landercasper, J., Cogbill, T.H., Strutt, P.J., Landercasper, B.O., 1988. Trauma and the veterinarian. *J. Trauma* 28, 1255–1258.
- Le Meur, A., Albert, J.-D., Gautier, P., Ropars, M., Perdriger, A., Gando, N., Minet, J., Cormier, M., Bonnaure-Mallet, M., Jolivet-Gougeon, A., 2008. Acute tenosynovitis of the ankle due to *Capnocytophaga cynodegmi*/*canimorsus* as identified by 16S rRNA gene sequencing. *Joint Bone Spine*, doi:10.1016/j.jbspin.2008.04.007, (Epub ahead of publication).
- LeMoal, G., Landron, C., Grollier, G., Robert, R., Burucoa, C., 2003. Meningitis due to *Capnocytophaga canimorsus* after receipt of a dog bite: case report and review of the literature. *Clin. Infect. Dis.* 36, e42–e46.
- Lion, C., Escande, F., Burdin, J.C., 1996. *Capnocytophaga canimorsus* infections in human: Review of the literature and cases report. *Eur. J. Epidemiol.* 12, 521–533.
- Macrea, M.M., McNamee, M., Martin, T.J., 2008. Acute onset of fever, chills and lethargy in a 36-year-old woman. *Chest* 133, 1505–1507.
- Mahrer, S., Raik, E., 1992. *Capnocytophaga canimorsus* septicemia associated with cat scratch. *Pathology* 24, 194–196.
- Mally, M., Cornelis, G.R., 2008. Genetic tools for *Capnocytophaga canimorsus*. *Appl. Environ. Microbiol.* 74, 6369–6377.
- Mally, M., Shin, H., Paroz, C., Landmann, R., Cornelis, G.R., 2008. *Capnocytophaga canimorsus*: a human pathogen feeding at the surface of epithelial cells and phagocytes. *PLoS Pathogens* 4, 1–11.
- Martone, M.J., Zuehl, R.W., Minson, G.E., Scheld, W.M., 1980. Post-splenectomy sepsis with DF-2: report of a case with isolation of the organism from the patient's dog. *Ann. Int. Med.* 93, 457–458.
- Matter, H.C., Sentinella, A., 1998. The epidemiology of bite and scratch injuries by vertebrate animals in Switzerland. *Eur. J. Epidemiol.* 14 (26), 483–490.
- McLean, C.R., Hargrove, R., Behn, E., 2004. The first fatal case of *Capnocytophaga canimorsus* sepsis caused by a cat scratch. *J. R. Nav. Serv.* 90, 13–15.
- Meybeck, A., Aoun, N., Granados, D., Pease, S., Yeni, P., 2006. Meningitis due to *Capnocytophaga canimorsus*: contribution of 16S RNA ribosomal sequencing for species identification. *Scand. J. Infect. Dis.* 38, 375–399.
- Meyer, S., Shin, H., Cornelis, G.R., 2008. *Capnocytophaga canimorsus* resists phagocytosis by macrophages and blocks the ability of macrophages to kill other bacteria. *Immunobiology* 213, 805–814.
- Meyers, B., Schoeman, J.P., Goddard, A., Picard, J., 2008. The bacteriology and antimicrobial susceptibility of infected and non-infected dog bite wounds: fifty cases. *Vet. Microbiol.* 127, 360–368.
- Mirza, I., Wolk, J., Toth, L., Rostenburg, P., Kranwinkel, R., Sieber, S.C., 2000. Waterhouse-Friderichsen syndrome secondary to *Capnocytophaga canimorsus* septicemia and demonstration of bacteria by peripheral blood smear. *Arch. Pathol. Lab. Med.* 124, 859–863.
- Mulder, A.H., Gerlag, P.G., Verhoef, L.H., van den Wall Bake, A.W., 2001. Hemolytic uremic syndrome after *Capnocytophaga canimorsus* (DF-2) septicemia. *Clin. Nephrol.* 44, 167–170.
- Newton, L., 2006. An unusual case of septicaemia. *N. Z. J. Med. Lab. Sci.* 60, 59–60.
- Ndon, J., 1992. *Capnocytophaga canimorsus* septicaemia caused by a dog bite in a hairy cell leukemia patient. *J. Clin. Microbiol.* 30, 211–213.
- Overall, K.L., Love, M., 2001. Dog bites to humans—demography, epidemiology, injury, and risk. *J. Am. Vet. Med. Assoc.* 218, 1923–1934.
- Papadaki, T.G., el Moussoui, R., van Ketel, R.J., Verbraak, F.D., Tan, H.S., 2008. *Capnocytophaga canimorsus* endogenous endophthalmitis in an immunocompetent host. *Br. J. Ophthalmol.* 92, 1566–1567.
- Pedersen, G., Schönheyder, H.C., Nielsen, L.C., 1993. *Capnocytophaga canimorsus* bacteraemia demonstrated by appositional blood smear: a case report. *Acta Pathol. Microbiol. Immunol. Scand.* 101, 572–574.
- Pers, C., Gahrn-Hansen, B., Frederiksen, W., 1996. *Capnocytophaga canimorsus* septicemia in Denmark 1982–1995: review of 39 cases. *Clin. Infect. Dis.* 23, 71–75.
- Phipps, S.E., Tamblyn, D.M., Badenoch, P.R., 2002. *Capnocytophaga canimorsus* endophthalmitis following cataract surgery. *Clin. Exp. Ophthalmol.* 30, 375–377.
- Piccininno, G., Palliola, E., Sala, V., 1984. Clinical and anatomohistological features of experimental infection by DF-2 in rabbits. *Path. Res. Pract.* 179, 95–100.
- Pletschette, M., Köhl, J., Kuipers, J., Schmidt, R.E., 1992. Opportunistic *Capnocytophaga canimorsus* infection. *Lancet* 339, 308.
- Risi, G.F., Spangler, C.A., 2006. *Capnocytophaga canimorsus* meningitis after routine myelography: a sentinel event identifies multiple opportunities for improvement of standard practices in radiology. *Am. J. Infect. Control* 34, 540–542.
- Roscoe, D.L., Zemcov, S.J.V., Thornber, D., Wise, R., Clarke, A.M., 1992. Antimicrobial susceptibilities and β -lactamase characterization of *Capnocytophaga* species. *Antimicrob. Agents Chemother.* 36, 2197–2200.
- Sandoe, J.A.T., 2004. *Capnocytophaga canimorsus* endocarditis. *J. Med. Microbiol.* 53, 245–248.
- Sawmiller, C., Dudrick, S.J., Hamzi, M., 1998. Postsplenectomy *Capnocytophaga canimorsus* sepsis presenting as an acute abdomen. *Arch. Surg.* 133, 1362–1365.
- Scarlett, J.D., Williamson, H.G., Dadson, P.J., Fassett, R., Peel, M., 1991. A syndrome resembling thrombotic thrombocytopenic purpura associated with *Capnocytophaga canimorsus* septicemia. *Am. J. Med.* 90, 127–128.
- Shin, H., Mally, M., Kuhn, M., Paroz, C., Cornelis, G.R., 2007. Escape from immune surveillance by *Capnocytophaga canimorsus*. *J. Infect. Dis.* 195, 375–386.
- Singer, D.B., 1973. Post-splenectomy sepsis. In: Rosenberg, H.S., Bolande, R.P. (Eds.), *Perspectives in Pediatric Pathology*, vol. 1. Year Book Medical Publishers, Chicago, IL, pp. 549–563.
- de Smet, M.D., Chan, C.C., Nussenblatt, R.B., Palestine, A.G., 1990. *Capnocytophaga canimorsus* as the cause of a chronic corneal infection. *Am. J. Ophthalmol.* 109, 240–242.
- Sowden, D., Allworth, A., Davis, L.B., 1995. *Capnocytophaga canimorsus* sepsis: poor growth using an automated blood culture system. *Clin. Microbiol. Newslett.* 17, 94–96.
- Speck, H., Kroppenstedt, R.M., Mannheim, W., 1987. Genomic relationships and species differentiation in the genus *Capnocytophaga*. *Zentralbl. Bakteriol. Mikrobiol. Hyg. (A)* 266, 390–402.
- Sumaraju, V., Smith, L.G., Smith, S.M., 2001. Infectious complications in asplenic hosts. *Infect. Dis. Clin. North Am.* 15, 551–565.
- Talan, D.A., Citron, D.M., Abrahamian, F.M., Gregory, D.M., Moran, J., Goldstein, E.J.C., 1999. Bacteriologic analysis of infected dog and cat bites. *N. Engl. J. Med.* 340, 85–92.
- Tierney, D.M., Strauss, L.P., Sanchez, J.L., 2006. *Capnocytophaga canimorsus* mycotic abdominal aortic aneurysm: why the mailman is afraid of dogs. *J. Clin. Microbiol.* 44, 649–651.
- Tison, D.L., Latimer, J.M., 1986. Lysis-centrifugation-direct plating technique for isolation of Group DF-2 from the blood of a dog bite victim. *J. Infect. Dis.* 153, 1001–1002.
- Tobé, T.J., Franssen, C.F., Zijlstra, J.G., de Jong, P.E., Stegeman, C.A., 1999. Hemolytic uremic syndrome due to *Capnocytophaga canimorsus* bacteraemia after a dog bite. *Am. J. Kidney Dis.* 33, e5.
- Underman, A.E., 1987. Bite wounds inflicted by dogs and cats. *Vet. Clin. North Am.* 17, 195–207.
- Valtonen, M., Lauhio, A., Carlson, P., Multanen, J., Sivonen, A., Vaara, M., Lähdevirta, J., 1995. *Capnocytophaga canimorsus* septicemia: fifth report of a cat-associated infection and five other cases. *Eur. J. Clin. Microbiol. Infect. Dis.* 14, 520–523.
- van de Ven, A.R.T., van Vliet, A.C.M., Maraha, B., Ponsen, H.H., 2004. Fibrinolytic therapy in *Capnocytophaga canimorsus* sepsis after dog bite. *Intensive Care Med.* 30, 1980.
- Verghese, A., Hamati, F., Berk, S., Franzus, B., Berk, S., Smith, J.K., 1988. Susceptibility of dysgonic fermenter 2 to antimicrobial agents in vitro. *Antimicrob. Agents Chemother.* 32, 78–80.
- Wareham, D.W., Michael, J.S., Warwick, S., Whitlock, P., Wood, A., Das, S.S., 2007. The dangers of dog bites. *J. Clin. Pathol.* 60, 328–329.
- Weinberg, E.D., 2000. Microbial pathogens with impaired ability to acquire host iron. *BioMetals* 13, 85–89.
- Westwell, A.J., Kerr, K., Spencer, M.B., Hutchinson, D.N., 1989. DF-2 infection. *Brit. Med. J.* 298, 116–117.
- Zimmer-Gallen, I.E., Pach, J.M., 1996. *Capnocytophaga canimorsus* endophthalmitis. *Retina* 16, 163–164.